

Petrified Trees – Their Formation Conditions, Types, and Distribution

Mari Kochalidze

mari.kochalidze511@hum.tsu.edu.ge

Department of Geology, Faculty of Exact and Natural Sciences, Ivane Javakhishvili Tbilisi State University, 13 University St., Tbilisi, Georgia, 0186

Petrified wood is formed as a result of the complete replacement of the organic material of wood by mineral matter, with full preservation of its internal structure. This process involves the phenomenon of pseudomorphism.

Petrified wood forms under specific environmental conditions. An oxygen-deficient environment and the presence of various mineralized solutions are essential. The replacement of organic material the “petrification” of wood is a long-term, multistage process that depends on the composition of the solution, as well as changes in temperature, pressure, and pH. Mineral replacement of the wood proceeds slowly and gradually, ensuring the preservation of its original structure.

Petrified trees have different mineral compositions and therefore display a variety of colors. Common types include silicified (quartzified), opalized, chalcedony-bearing, and other forms of petrified wood.

Petrified trees are an important object of study in geology and paleobotany. They provide valuable information about past ecosystems, climatic conditions, and geological processes. Petrified forests are distributed across various continents. They are protected as natural heritage sites and are used for scientific, educational, and tourism purposes.

This presentation discusses the formation conditions of petrified trees, their types, distribution, and significance. Special attention is given to the Goderdzi Petrified Forest, located in the Kvabliani River basin, the Dzindze Valley, on the eastern slope of the Arsiani Range, within the Adigeni and Khulo municipalities.